

Note to readers with disabilities: *EHP* strives to ensure that all journal content is accessible to all readers. However, some figures and Supplemental Material published in *EHP* articles may not conform to [508 standards](#) due to the complexity of the information being presented. If you need assistance accessing journal content, please contact ehp508@niehs.nih.gov. Our staff will work with you to assess and meet your accessibility needs within 3 working days.

Supplemental Material

Exposure Disparities by Income, Race and Ethnicity, and Historic Redlining Grade in the Greater Seattle Area for Ultrafine Particles and Other Air Pollutants

Kaya Bramble, Magali N. Blanco, Annie Doubleday, Amanda J. Gasset, Anjum Hajat, Julian D. Marshall, and Lianne Sheppard

Table of Contents

Table S1. Population-weighted mean pollutant concentration (2019-2020) in blocks by historical Home Owners' Loan Corporation (HOLC) redlining grade in Seattle when all blocks within the HOLC-graded region are included compared to when only blocks that received one grade are included.

Table S2. Population-weighted mean pollutant concentration (2019-2020) in blocks by historical Home Owners' Loan Corporation (HOLC) redlining grade in Seattle when blocks falling within multiple HOLC grades are split evenly by population compared to when the population split is weighted by area.

Table S3. Mean difference (95% CI) in pollutant concentration (2019-2020) per 10% increase in income (2006-2010 American Community Survey) in greater Seattle area from regression models with log-transformed (base 10) block level household income as predictor variable. Models accounted for spatial correlation using generalized estimating equations with identity link.

Table S4. Mean difference (95% CI) in pollutant concentration (2019-2020) per 5% increase in race and ethnicity group (2010 Census). Each group was modeled separately in population-weighted linear regression models with the percentage of that specific racial or ethnic group in the block as the predictor. Models accounted for spatial correlation using generalized estimating equations with identity link. This table provides the data for Figure 3.

Table S5. Population-weighted mean pollutant concentration (2019-2020) by block level income group (2006-2020 American Community Survey) restricted to the Home Owners' Loan Corporation (HOLC) redlining region in Seattle and the percent difference in comparison to the population-weighted average of the overall population in the HOLC-graded region.

Table S6. Population-weighted mean pollutant concentration (2019-2020) by racial and ethnic group (2010 Census) restricted to the Home Owners' Loan Corporation (HOLC) redlining region in Seattle and the percent difference in comparison to the population-weighted average of the overall population in the HOLC-graded region.

Figure S1. Study region with Seattle redlining Home Owner's Loan Corporation (HOLC) grades, from A ("best") to D ("hazardous") and X ("interior uncategorized"). Areas within the HOLC-graded region that do not show any blocks are areas without residential population in the 2010 Census. Maps generated in Quantum Geographic Information System (QGIS) using data from the American Community Survey (ACS), Census, and the University of Richmond's Mapping Inequality.

Figure S2. Scatterplots comparing log (base 10) of block group-level median household income to pollutant concentrations, with least squares regression line (red) and smoothing line (blue). Smoothing lines were generated using the `geom.smooth()` function in R.

Figure S3. Scatterplots comparing block-level percent Non-Hispanic residents to pollutant concentrations, with least squares regression line (red) and smoothing line (blue). Smoothing lines were generated using the `geom.smooth()` function in R.

Figure S4. Scatterplots comparing block-level percent Hispanic residents to pollutant concentrations, with least squares regression line (red) and smoothing line (blue). Smoothing lines were generated using the `geom.smooth()` function in R.

Figure S5. Scatterplots comparing block-level percent Black residents to pollutant concentrations, with least squares regression line (red) and smoothing line (blue). Smoothing lines were generated using the `geom.smooth()` function in R.

Figure S6. Scatterplots comparing block-level percent Asian residents to pollutant concentrations, with least squares regression line (red) and smoothing line (blue). Smoothing lines were generated using the `geom.smooth()` function in R.